arXiv:1202.2207v1 [math.CA] 10 Feb 2012

NEW ESTIMATIONS FOR $h-$CONVEX FUNCTIONS VIA FURTHER PROPERTIES

MEVLÜT TUNÇ♣ AND HÜSEYIN YILDIRIM♠

Abstract. In this paper, some new inequalities of the Hermite-Hadamard type for $h-$ convex functions whose modulus of the derivatives are h-convex and applications for special means are given.

1. Introduction

The following definition is well known in the literature [11]: A function $f : I \to \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$, is said to be convex on I if inequality

$$f\left(tx+(1-t)y\right) \leq tf\left(x\right)+(1-t)f\left(y\right) \tag{1.1}$$

holds for all $x, y \in I$ and $t \in [0,1]$. Geometrically, this means that if P, Q and R are three distinct points on the graph of f with Q between P and R, then Q is on or below chord PR.

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function and $a, b \in I$ with $a < b$. The following double inequality:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f\left(x\right)dx \leq \frac{f\left(a\right)+f\left(b\right)}{2} \tag{1.2}$$

is known in the literature as Hadamard's inequality (or H-H inequality) for convex function. Keep in mind that some of the classical inequalities for means can come from (1.2) for convenient particular selections of the function f. If f is concave, this double inequality hold in the inversed way.

Definition 1. *[8] We say that $f : I \to \mathbb{R}$ is Godunova-Levin function or that f belongs to the class $Q\left(I\right)$ if f is non-negative and for all $x, y \in I$ and $t \in (0,1)$ we have*

$$f\left(tx+(1-t)y\right) \leq \frac{f\left(x\right)}{t}+\frac{f\left(y\right)}{1-t}. \tag{1.3}$$

Definition 2. *[6] We say that $f : I \subseteq \mathbb{R} \to \mathbb{R}$ is a $P-$function or that f belongs to the class $P\left(I\right)$ if f is nonnegative and for all $x, y \in I$ and $t \in [0,1]$, we have*

$$f\left(tx+(1-t)y\right) \leq f\left(x\right)+f\left(y\right). \tag{1.4}$$

2000 *Mathematics Subject Classification.* 26A51, 26D10, 26D15.

Key words and phrases. $h-$convex function, Hermite-Hadamard inequality, Hölder inequality, Power-mean inequality, Special means.

♣Corresponding Author.

Definition 3. *[9] Let $s \in (0,1]$. A function $f : (0,\infty] \to [0,\infty]$ is said to be $s-$convex in the second sense if*

$$f(tx + (1-t)y) \le t^s f(x) + (1-t)^s f(y), \tag{1.5}$$

for all $x, y \in (0,b]$ and $t \in [0,1]$. This class of $s-$convex functions is usually denoted by K_s^2.

In 1978, Breckner introduced $s-$convex functions as a generalization of convex functions in [3]. Also, in that work Breckner proved the important fact that the set valued map is $s-$convex only if the associated support function is $s-$convex function in [4]. A number of properties and connections with s-convex in the first sense are discussed in paper [9]. Of course, $s-$convexity means just convexity when $s = 1$.

Definition 4. *[14] Let $h : J \subseteq \mathbb{R} \to \mathbb{R}$ be a positive function . We say that $f : I \subseteq \mathbb{R} \to \mathbb{R}$ is $h-$convex function, or that f belongs to the class $SX(h, I)$, if f is nonnegative and for all $x, y \in I$ and $t \in [0,1]$ we have*

$$f(tx + (1-t)y) \le h(t) f(x) + h(1-t) f(y). \tag{1.6}$$

If inequality (1.6) is reversed, then f is said to be $h-$concave, i.e. $f \in SV(h, I)$. Obviously, if $h(t) = t$, then all nonnegative convex functions belong to $SX(h, I)$ and all nonnegative concave functions belong to $SV(h, I)$; if $h(t) = \frac{1}{t}$, then $SX(h, I) = Q(I)$; if $h(t) = 1$, then $SX(h, I) \supseteq P(I)$; and if $h(t) = t^s$, where $s \in (0,1)$, then $SX(h, I) \supseteq K_s^2$.

Remark 1. *[14]Let h be a non-negative function such that*

$$h(\alpha) \ge \alpha \tag{1.7}$$

for all $\alpha \in (0,1)$. For example, the function $h_k(x) = x^k$ where $k \le 1$ and $x > 0$ has that property. If f is a non-negative convex function on I , then for $x, y \in I$, $\alpha \in (0,1)$ we have

$$f(\alpha x + (1-\alpha)y) \le \alpha f(x) + (1-\alpha) f(y) \le h(\alpha) f(x) + h(1-\alpha) f(y). \tag{1.8}$$

So, $f \in SX(h, I)$. Similarly, if the function h has the property: $h(\alpha) \le \alpha$ for all $\alpha \in (0,1)$, then any non-negative concave function f belongs to the class $SV(h, I)$.

For recent results and generalizations concerning $h-$convex functions see [2, 5, 12-14] and references therein.

In [7], the following theorem which was obtained by Dragomir and Agarwal contains the Hermite-Hadamard type integral inequality.

Theorem 1. *[7] Let $f : I^\circ \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I°, $a, b \in I^\circ$ with $a < b$. If $|f'|$ is convex on $[a,b]$, then the following inequality holds:*

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(u)\,du \right| \le \frac{(b-a)\left(|f'(a)| + |f'(b)|\right)}{8}. \tag{1.9}$$

In [10] Kavurmaci et. al. used the following lemma and introduced some new Hermite-Hadamard type inequalities:

Lemma 1. [10]*Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I°, where $a, b \in I$ with $a < b$. If $f' \in L[a,b]$, then the following equality holds:*

$$\begin{aligned} & \frac{(b-x) f(b) + (x-a) f(a)}{b-a} - \frac{1}{b-a} \int_a^b f(u)\, du \qquad (1.10) \\ = & \frac{(x-a)^2}{b-a} \int_0^1 (t-1) f'(tx + (1-t)a)\, dt \\ & + \frac{(b-x)^2}{b-a} \int_0^1 (1-t) f'(tx + (1-t)b)\, dt. \end{aligned}$$

Theorem 2. [10] *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I° sucht hat $f' \in L[a,b]$, where $a, b \in I$ with $a < b$. If $|f'|^{\frac{p}{p-1}}$ is convex on $[a,b]$ and for some fixed $q > 1$, then the following inequality holds:*

$$\begin{aligned} & \left| \frac{(b-x) f(b) + (x-a) f(a)}{b-a} - \frac{1}{b-a} \int_a^b f(u)\, du \right| \qquad (1.11) \\ \leq & \left(\frac{1}{1+p} \right)^{\frac{1}{p}} \left(\frac{1}{2} \right)^{\frac{1}{p}} \\ & \times \left[\frac{(x-a)^2 \left[|f'(a)|^q + |f'(x)|^q \right]^{\frac{1}{q}} + (b-x)^2 \left[|f'(x)|^q + |f'(b)|^q \right]^{\frac{1}{q}}}{b-a} \right] \end{aligned}$$

for each $x \in [a,b]$ and $q = \frac{p}{p-1}$.

In [1], Barani et. al. proved a variant of Hadamard's inequality which holds for $s-$convex functions in the second sense.

Theorem 3. [1] *Let $f : I \subseteq \mathbb{R}_+ \to \mathbb{R}$ be a differentiable mapping on I° and $a, b \in I^{\circ}$ with $a < b$. If $|f'|^{\frac{p}{p-1}}$ is $s-$convex on $[a,b]$ in the second sense, for some fixed $s \in (0,1]$. Then the following inequality holds,*

$$\begin{aligned} & \left| \frac{(b-x) f(b) + (x-a) f(a)}{b-a} - \frac{1}{b-a} \int_a^b f(u)\, du \right| \qquad (1.12) \\ \leq & \left(\frac{1}{1+p} \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left\{ \frac{(x-a)^2}{b-a} \left(|f'(x)|^q + \left[|f'(a)|^q \right]^{\frac{1}{q}} \right) \right. \\ & \left. + \frac{(b-x)^2}{b-a} \left(|f'(x)|^q + \left[|f'(b)|^q \right]^{\frac{1}{q}} \right) \right\}, \end{aligned}$$

where $q = \frac{p}{p-1}$.

The main purpose of this paper is to establish refinements inequalities of the results in [10]. We obtained new inequalities related to the right-hand side of Hermite-Hadamard inequality for functions when a power of whose first derivatives in absolute values is $h-$convex. Then, we give some applications for special means of real numbers.

2. Main Results

In this section we introduce some Hermite-Hadamard type inequalities for h-convex functions with corollaries and remarks.

Theorem 4. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I°, where $a, b \in I$ with $a < b$ such that $f' \in L[a,b]$ and h is supermultiplicative and nonnegative such that $h(\alpha) \geq \alpha$. If $|f'|$ is $h-$convex on $[a,b]$, then*

$$
\begin{aligned}
&\left|\frac{(b-x) f(b) + (x-a) f(a)}{b-a} - \frac{1}{b-a}\int_a^b f(u)\, du\right| \\
\leq\ & \frac{(x-a)^2}{b-a}\left(|f'(x)|\int_0^1 h((1-t)t)\, dt + |f'(a)|\int_0^1 h\left((1-t)^2\right) dt\right) \\
&+\frac{(b-x)^2}{b-a}\left(|f'(x)|\int_0^1 h((1-t)t)\, dt + |f'(b)|\int_0^1 h\left((1-t)^2\right) dt\right)
\end{aligned}
\tag{2.1}
$$

Proof. From Lemma 1, we have

$$
\begin{aligned}
&\left|\frac{(b-x) f(b) + (x-a) f(a)}{b-a} - \frac{1}{b-a}\int_a^b f(u)\, du\right| \\
=\ & \left|\frac{(x-a)^2}{b-a}\int_0^1 (t-1) f'(tx + (1-t)a)\, dt \right.\\
&\left. +\frac{(b-x)^2}{b-a}\int_0^1 (1-t) f'(tx + (1-t)b)\, dt\right| \\
\leq\ & \frac{(x-a)^2}{b-a}\int_0^1 (1-t)\,|f'(tx + (1-t)a)|\, dt \\
&+\frac{(b-x)^2}{b-a}\int_0^1 (1-t)\,|f'(tx + (1-t)b)|\, dt
\end{aligned}
$$

Since $|f'|$ is $h-$convex, then we obtain

$$
\begin{aligned}
&\left|\frac{(b-x) f(b) + (x-a) f(a)}{b-a} - \frac{1}{b-a}\int_a^b f(u)\, du\right| \\
\leq\ & \frac{(x-a)^2}{b-a}\int_0^1 (1-t)\left(h(t)\,|f'(x)| + h(1-t)\,|f'(a)|\right) dt \\
&+\frac{(b-x)^2}{b-a}\int_0^1 (1-t)\left(h(t)\,|f'(x)| + h(1-t)\,|f'(b)|\right) dt \\
\leq\ & \frac{(x-a)^2}{b-a}\left(|f'(x)|\int_0^1 h(1-t)\,h(t)\, dt + |f'(a)|\int_0^1 h^2(1-t)\, dt\right) \\
&+\frac{(b-x)^2}{b-a}\left(|f'(x)|\int_0^1 h(1-t)\,h(t)\, dt + |f'(b)|\int_0^1 h^2(1-t)\, dt\right) \\
\leq\ & \frac{(x-a)^2}{b-a}\left(|f'(x)|\int_0^1 h((1-t)t)\, dt + |f'(a)|\int_0^1 h\left((1-t)^2\right) dt\right) \\
&+\frac{(b-x)^2}{b-a}\left(|f'(x)|\int_0^1 h((1-t)t)\, dt + |f'(b)|\int_0^1 h\left((1-t)^2\right) dt\right)
\end{aligned}
$$

which completes the proof. □

Corollary 1. *In Theorem 4, if we choose* $x=\frac{a+b}{2}$, *we get*

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(u)\,du\right|\\ \leq\ &\frac{b-a}{4}\left(2\left|f'\left(\frac{a+b}{2}\right)\right|\int_0^1 h((1-t)t)\,dt+(|f'(a)|+|f'(b)|)\int_0^1 h\left((1-t)^2\right)dt\right)\end{aligned}$$

Corollary 2. *In Corollary 1, using the* $h-$*convexity of* $|f'|$, *we have*

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(u)\,du\right|\\ \leq\ &\frac{b-a}{4}(|f'(a)|+|f'(b)|)\left(2h\left(\frac{1}{2}\right)\int_0^1 h((1-t)t)\,dt+\int_0^1 h\left((1-t)^2\right)dt\right).\end{aligned}$$

Corollary 3. *In Corollary 2, if we choose* $h(t)=t$, *we have*

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(u)\,du\right|\\ \leq\ &\frac{b-a}{4}(|f'(a)|+|f'(b)|)\left(\int_0^1(1-t)t\,dt+\int_0^1(1-t)^2\,dt\right)\\ =\ &\frac{b-a}{8}(|f'(a)|+|f'(b)|)\end{aligned}$$

which is the inequality in (1.9).

Theorem 5. *Let* $f:I\subseteq\mathbb{R}\to\mathbb{R}$ *be a differentiable mapping on* I°, *where* $a,b\in I$ *with* $a<b$, *such that* $f'\in L[a,b]$ *and* $h:I\subseteq\mathbb{R}\to\mathbb{R}$ *be a nonnegative such that* $h\in L[0,1]$. *If* $|f'|^{\frac{p}{p-1}}$ *is* $h-$*convex on* $[a,b]$ *and for some fixed* $q>1$, *then*

$$\begin{aligned}&\left|\frac{(b-x)f(b)+(x-a)f(a)}{b-a}-\frac{1}{b-a}\int_a^b f(u)\,du\right|\\ \leq\ &\left(\frac{1}{1+p}\right)^{\frac{1}{p}}\left\{\frac{(x-a)^2}{b-a}\left(|f'(x)|^q\int_0^1 h(t)\,dt+|f'(a)|^q\int_0^1 h(1-t)\,dt\right)^{\frac{1}{q}}\right.\\ &\left.+\frac{(b-x)^2}{b-a}\left(|f'(x)|^q\int_0^1 h(t)\,dt+|f'(b)|^q\int_0^1 h(1-t)\,dt\right)^{\frac{1}{q}}\right\}\end{aligned} \tag{2.2}$$

for each $x\in[a,b]$ *and* $q=\frac{p}{p-1}$.

Proof. From Lemma 1 and using the well-known Hölder integral inequality, we obtain

$$\begin{aligned}
&\left|\frac{(b-x)\,f(b)+(x-a)\,f(a)}{b-a}-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{(x-a)^2}{b-a}\int_0^1 (1-t)\left|f'(tx+(1-t)\,a)\right|dt \\
&+\frac{(b-x)^2}{b-a}\int_0^1 (1-t)\left|f'(tx+(1-t)\,b)\right|dt \\
\leq\ & \left(\int_0^1 (1-t)^p\,dt\right)^{\frac{1}{p}}\left\{\frac{(x-a)^2}{b-a}\left(\int_0^1 \left|f'(tx+(1-t)\,a)\right|^q dt\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^2}{b-a}\left(\int_0^1 \left|f'(tx+(1-t)\,b)\right|^q dt\right)^{\frac{1}{q}}\right\}
\end{aligned}$$

Hence, by $h-$convexity of $|f'|^q$, we have

$$\begin{aligned}
&\left|\frac{(b-x)\,f(b)+(x-a)\,f(a)}{b-a}-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \left(\frac{1}{1+p}\right)^{\frac{1}{p}}\left\{\frac{(x-a)^2}{b-a}\left(|f'(x)|^q\int_0^1 h(t)\,dt+|f'(a)|^q\int_0^1 h(1-t)\,dt\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^2}{b-a}\left(|f'(x)|^q\int_0^1 h(t)\,dt+|f'(b)|^q\int_0^1 h(1-t)\,dt\right)^{\frac{1}{q}}\right\}
\end{aligned}$$

which completes the proof. □

Corollary 4. *In Theorem 5, if we choose $h(t)=t$, inequality (2.2) is reduces to (1.11)*

Corollary 5. *In Theorem 5, if we choose $h(t)=t^s$, inequality (2.2) is reduces to (1.12)*

Corollary 6. *In Theorem 5, if we choose $h(t)=1$, then we obtain an integral inequality for $P-$functions*

$$\begin{aligned}
&\left|\frac{(b-x)\,f(b)+(x-a)\,f(a)}{b-a}-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \left(\frac{1}{1+p}\right)^{\frac{1}{p}}\left[\frac{(x-a)^2+(b-x)^2}{b-a}\left(|f'(x)|^q+|f'(a)|^q\right)^{\frac{1}{q}}\right].
\end{aligned}$$

Remark 2. ○*In Theorem 5, choosing $x=a$, we get*

$$\begin{aligned}
&\left|f(b)-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & (b-a)\left(\frac{1}{1+p}\right)^{\frac{1}{p}}\left(|f'(a)|^q\int_0^1 h(t)\,dt+|f'(b)|^q\int_0^1 h(1-t)\,dt\right)^{\frac{1}{q}}.
\end{aligned}$$

○*In Theorem 5, choosing $x=b$, we get*

$$\begin{aligned}&\left|f\left(a\right)-\frac{1}{b-a}\int_{a}^{b}f\left(u\right)du\right|\\ \leq\quad&\left(b-a\right)\left(\frac{1}{1+p}\right)^{\frac{1}{p}}\left(\left|f'\left(b\right)\right|^{q}\int_{0}^{1}h\left(t\right)dt+\left|f'\left(a\right)\right|^{q}\int_{0}^{1}h\left(1-t\right)dt\right)^{\frac{1}{q}}.\end{aligned}$$

$\circ$*In Theorem 5, choosing* $x=\frac{a+b}{2}$, *we get*

$$\begin{aligned}&\left|\frac{f\left(b\right)+f\left(a\right)}{2}-\frac{1}{b-a}\int_{a}^{b}f\left(u\right)du\right|\\ \leq\quad&\frac{\left(b-a\right)}{4}\left(\frac{1}{1+p}\right)^{\frac{1}{p}}\left\{\left(\left|f'\left(\frac{a+b}{2}\right)\right|^{q}\int_{0}^{1}h\left(t\right)dt+\left|f'\left(a\right)\right|^{q}\int_{0}^{1}h\left(1-t\right)dt\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\left|f'\left(\frac{a+b}{2}\right)\right|^{q}\int_{0}^{1}h\left(t\right)dt+\left|f'\left(b\right)\right|^{q}\int_{0}^{1}h\left(1-t\right)dt\right)^{\frac{1}{q}}\right\}.\end{aligned}$$

$\circ$*In Theorem 5, choosing* $x=\frac{a+b}{2}$ *and* $f'\left(\frac{a+b}{2}\right)=0$, *we get*

$$\begin{aligned}&\left|\frac{f\left(b\right)+f\left(a\right)}{2}-\frac{1}{b-a}\int_{a}^{b}f\left(u\right)du\right|\\ \leq\quad&\frac{\left(b-a\right)}{4}\left(\frac{1}{1+p}\right)^{\frac{1}{p}}\left[\left(\left|f'\left(a\right)\right|+\left|f'\left(b\right)\right|\right)\left(\int_{0}^{1}h\left(t\right)dt\right)^{\frac{1}{q}}\right].\end{aligned}$$

Theorem 6. *Let* $f:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ *be a differentiable mapping on* I°, *where* $a,b\in I$ *with* $a<b$, *such that* $f'\in L\left[a,b\right]$ *and* $h:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ *be a nonnegative and supermultiplicative such that* $h\in L\left[0,1\right]$ *and* $h\left(t\right)\geq t$. *If* $\left|f'\right|^{q}$ *is* $h-$*convex on* $\left[a,b\right]$ *and for some fixed* $q>1$, *then*

$$\begin{aligned}&\left|\frac{\left(b-x\right)f\left(b\right)+\left(x-a\right)f\left(a\right)}{b-a}-\frac{1}{b-a}\int_{a}^{b}f\left(u\right)du\right|\\ \leq\quad&\left(\int_{0}^{1}h\left(1-t\right)dt\right)^{1-\frac{1}{q}}\\ &\times\left\{\frac{\left(x-a\right)^{2}}{b-a}\left(\left|f'\left(x\right)\right|^{q}\int_{0}^{1}h\left(t-t^{2}\right)dt+\left|f'\left(a\right)\right|^{q}\int_{0}^{1}h\left(\left(1-t\right)^{2}\right)dt\right)^{\frac{1}{q}}\right.\\ &\left.+\frac{\left(b-x\right)^{2}}{b-a}\left(\left|f'\left(x\right)\right|^{q}\int_{0}^{1}h\left(t-t^{2}\right)dt+\left|f'\left(b\right)\right|^{q}\int_{0}^{1}h\left(\left(1-t\right)^{2}\right)dt\right)^{\frac{1}{q}}\right\}\end{aligned}$$

for each $x\in\left[a,b\right]$.

Proof. From Lemma 1 and using the power-mean inequality for $q \geq 1$ and $p = \frac{q}{q-1}$ we obtain

$$\begin{aligned}
&\left|\frac{(b-x)\,f(b)+(x-a)\,f(a)}{b-a}-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{(x-a)^2}{b-a}\int_0^1 (1-t)\left|f'(tx+(1-t)\,a)\right| dt \\
&+\frac{(b-x)^2}{b-a}\int_0^1 (1-t)\left|f'(tx+(1-t)\,b)\right| dt \\
\leq\ & \left(\int_0^1 (1-t)\,dt\right)^{1-\frac{1}{q}}\left\{\frac{(x-a)^2}{b-a}\left(\int_0^1 (1-t)\left|f'(tx+(1-t)\,a)\right|^q dt\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^2}{b-a}\left(\int_0^1 (1-t)\left|f'(tx+(1-t)\,b)\right|^q dt\right)^{\frac{1}{q}}\right\}
\end{aligned}$$

Hence, by $h-$convexity of $|f'|^q$, we get

$$\begin{aligned}
&\left|\frac{(b-x)\,f(b)+(x-a)\,f(a)}{b-a}-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \left(\int_0^1 (1-t)\,dt\right)^{1-\frac{1}{q}}\left\{\frac{(x-a)^2}{b-a}\left(\int_0^1 (1-t)\left|f'(tx+(1-t)\,a)\right|^q dt\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^2}{b-a}\left(\int_0^1 (1-t)\left|f'(tx+(1-t)\,b)\right|^q dt\right)^{\frac{1}{q}}\right\} \\
\leq\ & \left(\int_0^1 h(1-t)\,dt\right)^{1-\frac{1}{q}}\left\{\frac{(x-a)^2}{b-a}\left(\int_0^1 h(1-t)\left\{h(t)\left|f'(x)\right|^q+h(1-t)\left|f'(a)\right|^q\right\} dt\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^2}{b-a}\left(\int_0^1 h(1-t)\left\{h(t)\left|f'(x)\right|^q+h(1-t)\left|f'(b)\right|^q\right\} dt\right)^{\frac{1}{q}}\right\}
\end{aligned}$$

which completes the proof. □

Corollary 7. *In Theorem 6, choosing $x = (a+b)/2$ and then using the $h-$convexity of $|f'|^q$, we get*

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{b-a}{4}\left(\int_0^1 h(1-t)\,dt\right)^{\frac{1}{p}}\left\{\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q\int_0^1 h\left(t-t^2\right)dt+\left|f'(a)\right|^q\int_0^1 h\left((1-t)^2\right)dt\right)^{\frac{1}{q}}\right. \\
&\left.+\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q\int_0^1 h\left(t-t^2\right)dt+\left|f'(b)\right|^q\int_0^1 h\left((1-t)^2\right)dt\right)^{\frac{1}{q}}\right\}.
\end{aligned}$$

Remark 3. *In Corollary 7, choosing* $h(t)=t$*, we get*

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{b-a}{4}\left(\frac{1}{2}\right)^{\frac{1}{p}}\left(\frac{1}{3}\right)^{\frac{1}{q}}\left\{\left(\frac{1}{2}\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(a)|^q\right)^{\frac{1}{q}}\right. \\
&\left.+\left(\frac{1}{2}\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(b)|^q\right)^{\frac{1}{q}}\right\}
\end{aligned}$$

Corollary 8. *In Corollary 7, if we choose* $h(t)=1$*, then we obtain an integral inequality for* $P-$*functions*

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{b-a}{4}\left\{\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(a)|^q\right)^{\frac{1}{q}}\right. \\
&\left.+\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(b)|^q\right)^{\frac{1}{q}}\right\} \\
\leq\ & \frac{b-a}{2}\left(|f'(a)|+|f'(b)|\right).
\end{aligned}$$

3. Applications to Special Means

We now consider the applications of our Theorems to the following special means
The quadratic mean:

$$K=K(a,b):=\sqrt{\frac{a^2+b^2}{2}}\ a,b\geq 0,$$

The arithmetic mean:

$$A=A(a,b):=\frac{a+b}{2},\quad a,b\geq 0,$$

The geometric mean:

$$G=G(a,b):=\sqrt{ab},\ a,b\geq 0,$$

The logarithmic mean:

$$L=L(a,b):=\begin{cases} a & \text{if}\quad a=b \\ \frac{b-a}{\ln b-\ln a} & \text{if}\quad a\neq b\end{cases},\ a,b\geq 0$$

The p-logarithmic mean:

$$\begin{aligned}
L_p &= L_p(a,b):=\begin{cases}\left[\frac{b^{p+1}-a^{p+1}}{(p+1)(b-a)}\right]^{1/p} & \text{if}\quad a\neq b \\ a & \text{if}\quad a=b\end{cases}, \\
p &\in \mathbb{R}\backslash\{-1,0\};\ a,b>0.
\end{aligned}$$

Proposition 1. *Let $a, b \in \mathbb{R}$, $0 < a < b$ and $n \in \mathbb{Z}$, $|n| \geq 1$. Then, the following inequality holds,*

$$\left|A\left(a^{n}, b^{n}\right)-L_{n}^{n}(a, b)\right| \leq|n| \frac{b-a}{12}\left(A^{n-1}(a, b)+2 A\left(a^{n-1}, b^{n-1}\right)\right) . \tag{3.1}$$

Proof. If we apply Corollary 1 for $f(x)=x^{n}$, $h(t)=t$ where $x \in \mathbb{R}$, $n \in \mathbb{Z}$, $|n| \geq 1$, we get the proof (3.1). □

Proposition 2. *Let $a, b \in \mathbb{R}$, $0 < a < b$ and $n \in \mathbb{Z}$, $|n| \geq 1$. Then, we have:*

$$\left|A\left(a^{-1}, b^{-1}\right)-L^{-1}(a, b)\right| \leq \frac{b-a}{12}\left(\frac{1}{A^{2}(a, b)}+\frac{2 K^{2}(a, b)}{G^{4}(a, b)}\right) \tag{3.2}$$

Proof. The proof is immediate from Corollary 1 applied for $f(x)=\frac{1}{x}$, $x \in[a, b]$ and $h(t)=t$. □

Proposition 3. *Let $a, b \in \mathbb{R}$, $0 < a < b$ and $n \in \mathbb{Z}$, $|n| \geq 1$. Then, the following inequality holds,*

$$\left|A\left(a^{n}, b^{n}\right)-L_{n}^{n}(a, b)\right| \leq \frac{b-a}{4}|n| A\left(a^{n-1}, b^{n-1}\right) . \tag{3.3}$$

Proof. The assertion follows from Corollary 2 applied to $f(x)=x^{n}$, $h(t)=t$ where $x \in \mathbb{R}$, $n \in \mathbb{Z}$, $|n| \geq 1$. □

Proposition 4. *Let $a, b \in \mathbb{R}$, $0 < a < b$ and $n \in \mathbb{Z}$, $|n| \geq 1$. Then, we have:*

$$\left|A\left(a^{-1}, b^{-1}\right)-L^{-1}(a, b)\right| \leq \frac{b-a}{8} \frac{2 K^{2}(a, b)}{G^{4}(a, b)} \tag{3.4}$$

Proof. The assertion follows from Corollary 2 applied to $f(x)=\frac{1}{x}$, $x \in[a, b]$ and $h(t)=t$. □

Proposition 5. *Let $a, b \in \mathbb{R}$, $0 < a < b$ and $n \in \mathbb{Z}$, $|n| \geq 1$. Then, for all $q > 1$:*

$$\begin{aligned} & \left|A\left(a^{n}, b^{n}\right)-L_{n}^{n}(a, b)\right| \\ \leq\ & |n|(b-a)(2)^{\frac{1}{q}-3}(3)^{-\frac{1}{q}}\left\{\left(\frac{A^{q(n-1)}(a, b)}{2}+a^{q(n-1)}\right)^{\frac{1}{q}}\right. \\ & \left.+\left(\frac{A^{q(n-1)}(a, b)}{2}+b^{q(n-1)}\right)^{\frac{1}{q}}\right\} . \end{aligned} \tag{3.5}$$

Proof. If we apply Corollary 7 for $f(x)=x^{n}$ where $x \in \mathbb{R}$, $n \in \mathbb{Z}$, $|n| \geq 1$, we get the proof (3.5). □

References

[1] Barani, A., Ghazanfari, A.G. and Dragomir, S.S. *New Hermite-Hadamard Inequalities for $s-$Convex Functions*, RGMIA Research Report Collection Volume 14, Article 101, 2011.

[2] Bombardelli, M., Varošanec, S. *Properties of $h-$convex functions related to the Hermite–Hadamard–Fejér inequalities*, Comput. Math. Appl. 58 (2009) 1869–1877.

[3] Breckner, W.W. *Stetigkeitsaussagen f¨ur eine Klasse verallgemeinerter konvexer funktionen in topologischen linearen Raumen*, Pupl. Inst. Math., 23, 13–20, 1978.

[4] Breckner, W. W. *Continuity of generalized convex and generalized concave set-valued functions*, Rev Anal. Num´er. Thkor. Approx., 22, 39–51, 1993.

[5] Burai, P., Hazy, A. *On approximately $h-$convex functions,* J. Convex Anal. 18 (2) (2011) 447–454.

[6] Dragomir, S.S., Pečarić, J. and Persson, L.E. *Some inequalities of Hadamard type*, Soochow J.Math., 21, 335–341, 1995.

[7] Dragomir, SS, Agarwal, RP. *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula.* Appl Math Lett. 11(5), 91–95 (1998). doi:10.1016/S0893-9659(98)00086-X

[8] Godunova, E.K., Levin, V.I. *Neravenstva dlja funkcii sirokogo klassa, soderzascego vypuklye, monotonnye i nekotorye drugie vidy funkii*, Vycislitel. Mat. i. Fiz. Mezvuzov. Sb. Nauc. Trudov, MGPI,Moskva, pp. 138–142, 1985.

[9] Hudzik, H., Maligranda, L. *Some remarks on $s-$convex functions*, Aequationes Math., 48, 100–111, 1994.

[10] Kavurmaci, H., Avci, M. and Ozdemir, M Emin. *New inequalities of Hermite-Hadamard type for convex functions with applications*, J. Ineq. Appl., 86 2011,.

[11] Mitrinović, D.S., Pečarić, J. and Fink, A.M. *Classical and new inequalities in analysis*, KluwerAcademic, Dordrecht, 1993.

[12] Özdemir, M.E., Gürbüz, M. and Akdemir, A.O. *Inequalities for $h-$Convex Functions via Further Properties*, RGMIA Research Report Collection Volume 14, article 22, 2011.

[13] Sarıkaya, M.Z., Sağlam, A., Yıldırım, H. *On some Hadamard-type inequalities for $h-$convex functions,* J. Math. Inequal. 2 (3) (2008) 335–341.

[14] Varošanec, S. *On $h-$convexity*, J. Math. Anal. Appl., Volume 326, Issue 1, 303–311, 2007.

♣Kilis 7 Aralik University, Faculty of Science and Arts, Department of Mathematics, 79000, Kilis, TURKEY
E-mail address: `mevluttunc@kilis.edu.tr`

♠Kahramanmaraş Sütçü İmam University, Faculty of Science and Arts, Department of Mathematics, 46000, Kahramanmaraş, TURKEY
E-mail address: `hyildir@ksu.edu.tr`